# [TRANSCRIPT]

SUPER: Douglas. Diagnosed with type 2 diabetes 13 years ago

**Douglas:** My goals are to be healthy, and to make sure I'm doing my best to manage my diabetes. My diabetes has changed over time, and I wonder if the way I manage it needs to as well. I'm ready to talk to my doctor, and learn about my options.

#### SUPER:

Please see Full Important Safety Information and link to Full Prescribing Information for Toujeo<sup>®</sup> next to this video.

#### SUPER:

Vanessa. Diabetes Educator \*Terms and restrictions apply. See website for full offer terms and eligibility restrictions.

Please see Full Important Safety Information and link to Full Prescribing Information for Toujeo<sup>®</sup> next to this video.

Vanessa: One option is Toujeo®

#### SUPER:

Please see Full Important Safety Information and link to Full Prescribing Information for Toujeo<sup>®</sup> next to this video.

Narrator: Insulin glargine injection 300 Units/mL.

# SUPER:

\*Terms and restrictions apply. See website for full offer terms and eligibility restrictions.

# Pay as low as \$0 and no more than \$99\* for a 30-day supply.

Please see Full Important Safety Information and link to Full Prescribing Information for Toujeo<sup>®</sup> next to this video.

**Vanessa:** Toujeo<sup>®</sup> is a long-acting man-made insulin used to control high blood sugar in adults and children who are 6 years of age and older with diabetes mellitus.

- Toujeo® is not for use to treat diabetic ketoacidosis
- It is not known if Toujeo® is safe and effective in children under 6 years of age

# SUPER:

Prescription Toujeo<sup>®</sup> is a long-acting man-made insulin used to control high blood sugar in adults and children who are 6 years of age and older with diabetes mellitus.

- Toujeo® is not for use to treat diabetic ketoacidosis
- It is not known if Toujeo<sup>®</sup> is safe and effective in children under 6 years of age

Please see Full Important Safety Information and link to Full Prescribing Information

for Toujeo® next to this video.

**Vanessa:** Once-daily Toujeo<sup>®</sup> is proven to provide blood sugar control, and significant A1C reductions for adults with type 1 and type 2 diabetes.

# SUPER:

From the makers of Lantus<sup>®</sup> (insulin glargine injection) 100 Units/mL Toujeo<sup>®</sup> provides:

Proven ALL Day & Night Blood Sugar Control

Proven A1C reductions

#### SUPER:

Toujeo<sup>®</sup> should be taken at the same time once a day.

#### SUPER:

\*Terms and restrictions apply. See website for full offer terms and eligibility restrictions.

Pay as low as \$0 and no more than \$99\* for a 30-day supply.

Please see Full Important Safety Information and link to Full Prescribing Information for Toujeo® next to this video.

# SUPER:

Important Safety Information for Toujeo® (insulin glargine injection) 300 Units/mL

**Narrator:** Do not take Toujeo<sup>®</sup> if you have low blood sugar or if you are allergic to insulin or any of the ingredients in Toujeo<sup>®</sup>.

#### SUPER:

Do not take Toujeo<sup>®</sup> if you have low blood sugar or if you are allergic to insulin or any of the ingredients in Toujeo<sup>®</sup>.

Please see Full Important Safety Information and link to Full Prescribing Information for Toujeo<sup>®</sup> next to this video.

# SUPER:

Once-daily Toujeo® should be taken once a day at the same time each day

Please see Full Important Safety Information and link to Full Prescribing Information for Toujeo® next to this video.

**Vanessa:** Toujeo<sup>®</sup> works by slowly releasing small amounts of insulin to provide continuous glucose-lowering activity that lasts beyond 24 hours.

# SUPER:

Blood sugar-lowering activity continues around the clock

Once-daily Toujeo® should be taken once a day at the same time each day

Please see Full Important Safety Information and link to Full Prescribing Information for Toujeo<sup>®</sup> next to this video.

#### SUPER:

Blood sugar levels stayed within the ADA target zone (80-130 mg/dL) up to 36 hours

Results from this controlled study of 30 patients may not apply in everyday use. Remember your blood sugar levels may vary depending on your diet, physical activity, medications, and other factors.

\*American Diabetes Association. *Diabetes Care.* 2020;43 (suppl 1): S66-S76.

Range within target zone

Toujeo® should be taken once a day at the same time each day

Please see Full Important Safety Information and link to Full Prescribing Information for Toujeo® next to this video.

# What is Toujeo<sup>®</sup> (insulin glargine injection) 300 Units/mL?

Prescription Toujeo is a long-acting man-made insulin used to control high blood sugar in adults and children who are 6 years of age and older with diabetes mellitus.

- Toujeo is not for use to treat diabetic ketoacidosis
- It is not known if Toujeo is safe and effective in children under 6 years of age

# Important Safety Information

Do not take Toujeo if you have low blood sugar or if you are allergic to insulin or any of the ingredients in Toujeo.

**Do not share your pen(s) with other people, even if the needle has been changed**. You may give other people a serious infection, or get a serious infection from them.

Before starting Toujeo, tell your doctor about all your medical conditions, including if you have liver or kidney problems, if you are pregnant or planning to become pregnant, or if you are breastfeeding or planning to breastfeed.

**Change (rotate) your injection sites within the area you chose with each dose** to reduce your risk of getting lipodystrophy (pitted or thickened skin) and localized cutaneous amyloidosis (skin with lumps) at the injection sites. **Do not** use the same spot for each injection or inject where the skin is pitted, thickened, lumpy, tender, bruised, scaly, hard, scarred or damaged.

Heart failure can occur if you are taking insulin together with pills called TZDs (thiazolidinediones), even if you have never had heart failure or other heart problems. If you have heart failure, it may get worse while you take TZDs with Toujeo. Your treatment with TZDs and Toujeo may need to be changed or stopped by your doctor if you have new or worsening heart failure. Tell your doctor if you have any new or worsening symptoms including:

- Shortness of breath
- Sudden weight gain

• Swelling of your ankles or feet

Tell your doctor about all the medications you take, including OTC medicines, vitamins, and supplements, and herbal supplements.

Toujeo should be taken at the same time once a day. Test your blood sugar levels daily while using any insulin. Do not change your dose or type of insulin without talking to your doctor. Verify you have the correct insulin before each injection. **Do NOT use a syringe to remove Toujeo from your pen.** Your dose for Toujeo may be different from other insulins you have taken. Any change of insulin should be made cautiously and only under medical supervision.

**Do NOT dilute or mix Toujeo with any other insulin or solution**. It will not work as intended and you may lose blood sugar control, which could be serious. Use Toujeo only if the solution is clear and colorless with no particles visible.

While using Toujeo, do not drive or operate heavy machinery until you know how Toujeo affects you. Don't drink alcohol or use other medicines that contain alcohol.

The most common side effect of any insulin, including Toujeo, is low blood sugar (hypoglycemia), which may be serious and life-threatening. Severe hypoglycemia may cause harm to your heart or brain. Symptoms of serious low blood sugar may include shaking, sweating, fast heartbeat, and blurred vision.

Toujeo may cause severe allergic reactions that can lead to death. Get medical help right away if you have:

- A rash over your whole body
- Shortness of breath
- Swelling of your face, tongue, or throat
- Extreme drowsiness, dizziness, or confusion
- Trouble breathing
- Fast heartbeat
- Sweating

Toujeo may have additional side effects including swelling, weight gain, low potassium, and injection site reactions which may include change in fat tissue, skin thickening, redness, swelling, and itching.

Toujeo® SoloStar® and Toujeo® Max SoloStar® are single-patient-use prefilled insulin

pens. It is important to perform a safety test when using a new pen for the first time. Talk to your doctor about proper injection technique and follow instructions in the *Instruction Leaflet* that comes with the pen.

# SUPER:

# Please see full Prescribing Information for Toujeo® at link below

#### SUPER:

Please see Full Important Safety Information and link to Full Prescribing Information for Toujeo® next to this video.

SAUS.TJO.18.04.2722(1) | 01/20