

MAT-US-2007148 v2.0 Toujeo DTC Branded Doug & Vanessa Video Toujeo.com 2022 ISI
Update Transcript

From 00:00:01-00:00:13

My goals are to be healthy and make sure I'm doing my best to manage my diabetes. My diabetes has changed over time and I wonder if the way I manage it needs to as well, I'm ready to talk to my doctor and learn about my options.

From 00:00:14-00:00:20

One option is Toujeo (insulin glargine) injection 300 Units/mL.

From 00:00:21-00:00:39

Prescription Toujeo is a long-acting man-made insulin used to control high blood sugar in adults and children who are 6 years of age and older with diabetes mellitus.

- Toujeo is not for the treatment of diabetic ketoacidosis
- It is not known if Toujeo is safe and effective in children under 6 years of age.

From 00:00:40-00:00:49

once daily Toujeo is proven to provide blood sugar control and significant A1C reduction for adults with type 1 and type 2 diabetes.

From 00:00:50-00:00:55

Do not use Toujeo if you have low blood sugar or if you are allergic to insulin or any of the ingredients in Toujeo.

From 00:01:02-00:01:10

Toujeo works by slowly releasing small amounts of insulin to provide continuous glucose lowering activities that last beyond 24 hours.

From 00:01:21-00:05:11

What is Toujeo U-300 (insulin glargine) injection 300 Units/mL?

Prescription Toujeo is a long-acting man-made insulin used to control high blood sugar in adults and children who are 6 years of age and older with diabetes mellitus.

- Toujeo is not for the treatment of diabetic ketoacidosis
- It is not known if Toujeo is safe and effective in children under 6 years of age

Important Safety Information for Toujeo U-300 (insulin glargine) injection

Do not use Toujeo if you have low blood sugar or if you are allergic to insulin or any of the ingredients in Toujeo.

Do not share your pen(s) with other people, even if the needle has been changed. You may give other people a serious infection or get a serious infection from them.

Before starting Toujeo, tell your doctor about all your medical conditions, including if you have liver or kidney problems, if you are pregnant or planning to become pregnant, or if you are breastfeeding or planning to breastfeed.

Change (rotate) your injection sites within the area you chose with each dose to reduce your risk of getting pitted or thickened skin (lipodystrophy) and skin with lumps (localized cutaneous amyloidosis) at the injection sites. **Do not** use the same spot for each injection or inject where the skin is pitted, thickened, lumpy, tender, bruised, scaly, hard, scarred, or damaged.

Heart failure can occur if you are taking insulin together with pills called TZDs (thiazolidinediones), even if you have never had heart failure or other heart problems. If you have heart failure, it may get worse while you take TZDs with Toujeo. Your treatment with TZDs and Toujeo may need to be changed or stopped by your doctor if you have new or worsening heart failure. Tell your doctor if you have any new or worsening symptoms including:

- Shortness of breath
- Sudden weight gain
- Swelling of your ankles or feet

Tell your doctor about all the medications you take, including OTC medicines, vitamins, and supplements, and herbal supplements.

Toujeo should be taken at the same time once a day. Test your blood sugar levels daily while using any insulin. Do not change your dose or type of insulin without talking to your doctor. Verify you have the correct insulin before each injection. **Do NOT use a syringe to remove Toujeo from your pen.** Your dose for Toujeo may be different from other insulins you have taken. Any change of insulin should be made cautiously and only under medical supervision.

Do NOT dilute or mix Toujeo with any other insulin or solution. It will not work as intended and you may lose blood sugar control, which could be serious. Use Toujeo only if the solution is clear and colorless with no particles visible.

While using Toujeo, do not drive or operate heavy machinery until you know how Toujeo affects you. Don't drink alcohol or use other medicines that contain alcohol.

The most common side effect of Toujeo is low blood sugar (hypoglycemia), which may be serious and life-threatening. Severe hypoglycemia may cause harm to your heart or brain. Symptoms of serious low blood sugar may include shaking, sweating, fast heartbeat, and blurred vision. The long-acting effect of Toujeo may delay recovery from low blood sugar compared to shorter-acting insulins.

Toujeo may cause severe allergic reactions that can lead to death. Get medical help right away if you have:

- A rash over your whole body
- Shortness of breath
- Swelling of your face, tongue, or throat
- Extreme drowsiness, dizziness, or confusion
- Trouble breathing
- Fast heartbeat
- Sweating

Toujeo may have additional side effects including swelling, weight gain, low potassium, and injection site reactions which may include change in fat tissue, skin thickening, redness, swelling, and itching.

Toujeo SoloStar and Toujeo Max SoloStar are single-patient-use prefilled insulin pens. It is important to perform a safety test when using a new pen for the first time. Talk to your doctor about proper injection technique and follow instructions in the *Instruction Leaflet* that comes with your Toujeo SoloStar or Toujeo Max SoloStar pen.

©2022 sanofi-aventis U.S. LLC. All rights reserved.

MAT-US-2007148-v2.0-11/2022