

## TIME TO THINK ABOUT INSULIN? See how Toujeo Max can mean more.



As you and your doctor decide which insulin is right for you, ask your doctor if the Toujeo<sup>®</sup> Max SoloStar<sup>®</sup> pen is a possibility. Your insulin dose may change over time as your insulin needs change, and the Max SoloStar pen allows for a broad range of dosing. So you have the option of increasing your dose without changing your long-acting insulin pen. Depending on your daily dose and insurance coverage, using the Max SoloStar pen may mean fewer pens, fewer refills, and fewer associated co-pays. The Max SoloStar pen is recommended for patients who need at least 20 units of long-acting insulin per day.



### FEWER PENS

Toujeo is the most concentrated long-acting insulin with **the lowest injection volume**. Each unit of insulin is smaller, so you inject less liquid with your prescribed dose. That, plus a 900-unit pen, could mean you use fewer pens.



### FEWER REFILLS

The Max SoloStar pen holds more units of insulin than any other long-acting pen, so you may need to refill your prescription less often.




### FEWER CO-PAYS\*

If you're using fewer insulin pens, you may have fewer refills and associated co-pays, depending on daily dose and prescription insurance coverage.

[Find out more](#) ▶

\*Fewer co-pays does not guarantee savings.

### LOWEST INJECTION VOLUME COMPARED TO A UNIT OF ANY OTHER LONG-ACTING INSULIN\*



**WHEN YOU'RE READY TO START INSULIN, ASK YOUR DOCTOR  
IF YOU SHOULD START WITH THE TOUJEO MAX SOLOSTAR PEN.**

### What is Toujeo U-300 (insulin glargine) injection 300 Units/mL?

Prescription Toujeo is a long-acting man-made insulin used to control high blood sugar in adults and children who are 6 years of age and older with diabetes mellitus.

- Toujeo is not for the treatment of diabetic ketoacidosis
- It is not known if Toujeo is safe and effective in children under 6 years of age

### Important Safety Information for Toujeo U-300 (insulin glargine) injection

Do not use Toujeo if you have low blood sugar or if you are allergic to insulin or any of the ingredients in Toujeo.

Please see full Important Safety Information for Toujeo throughout. Please [click here](#) or visit <https://products.sanofi.us/toujeo/toujeo.pdf> for full Prescribing Information for Toujeo.

### QUESTIONS TO ASK YOURSELF BEFORE YOU VISIT YOUR DOCTOR:

1. What do I need to know about starting insulin? [Learn more](#) ▶

---

2. What are the signs that my diabetes may be changing? [Learn more](#) ▶

---

3. Why isn't it enough to take non-insulin medications? [Learn more](#) ▶

---

4. Could a long-acting insulin like Toujeo help me work toward my blood sugar and A1C goals?

---

5. What are the side effects of Toujeo? Is there something I can do to prepare?

---

### QUESTIONS TO ASK YOUR DOCTOR:

1. For my first insulin, is the 900-unit Toujeo<sup>®</sup> Max SoloStar<sup>®</sup> pen right for me?

---

2. Will my dose stay the same once I do start on insulin?

---

3. What is injecting with the Toujeo Max SoloStar pen like? [Check out this video](#) ▶

---

4. Does an insulin that is more concentrated mean I might have less liquid (volume) to inject? [Learn more](#) ▶

---

5. What if I skip or forget a dose of Toujeo?

---

### Important Safety Information for Toujeo U-300 (insulin glargine) injection

**Do not share your pen(s) with other people, even if the needle has been changed.** You may give other people a serious infection or get a serious infection from them.

Before starting Toujeo, tell your doctor about all your medical conditions, including if you have liver or kidney problems, if you are pregnant or planning to become pregnant, or if you are breastfeeding or planning to breastfeed.

Please see full Important Safety Information for Toujeo throughout. Please [click here](#) or visit <https://products.sanofi.us/toujeo/toujeo.pdf> for full Prescribing Information for Toujeo.

### QUESTIONS YOUR DOCTOR MAY HAVE FOR YOU:

1. Have you been tracking your blood sugar?
2. Have you experienced any symptoms?
3. Have you been following your treatment plan?

### NEED HELP ANSWERING THEM?

1. Bring your blood sugar log with you. Need a new one? [Get it here](#) ▶
2. Jot down any symptoms you've experienced—even if you start today.
3. Sticking to your plan helps your doctor know when changes need to be made.

### Are you prepared?

#### HERE'S WHAT TO BRING TO YOUR NEXT DOCTOR'S VISIT:

- |  |   |
|--|---|
| <input type="radio"/> Insurance card                                     | <input type="radio"/> Blood sugar log             |
| <input type="radio"/> Physician referral (if you're seeing a specialist) | <input type="radio"/> List of current medications |
| <input type="radio"/> Medical history                                    |   |



### Important Safety Information for Toujeo U-300 (insulin glargine) injection (cont'd)

**Change (rotate) your injection sites within the area you chose with each dose** to reduce your risk of getting pitted or thickened skin (lipodystrophy) and skin with lumps (localized cutaneous amyloidosis) at the injection sites. **Do not** use the same spot for each injection or inject where the skin is pitted, thickened, lumpy, tender, bruised, scaly, hard, scarred, or damaged.

Heart failure can occur if you are taking insulin together with pills called TZDs (thiazolidinediones), even if you have never had heart failure or other heart problems. If you have heart failure, it may get worse while you take TZDs with Toujeo. Your treatment with TZDs and Toujeo may need to be changed or stopped by your doctor if you have new or worsening heart failure. Tell your doctor if you have any new or worsening symptoms including:

- Shortness of breath
- Sudden weight gain
- Swelling of your ankles or feet

Tell your doctor about all the medications you take, including OTC medicines, vitamins, and supplements, and herbal supplements.

Toujeo should be taken at the same time once a day. Test your blood sugar levels daily while using any insulin. Do not change your dose or type of insulin without talking to your doctor. Verify you have the correct insulin before each injection. **Do NOT use a syringe to remove Toujeo from your pen.** Your dose for Toujeo may be different from other insulins you have taken. Any change of insulin should be made cautiously and only under medical supervision.

**Do NOT dilute or mix Toujeo with any other insulin or solution.** It will not work as intended and you may lose blood sugar control, which could be serious. Use Toujeo only if the solution is clear and colorless with no particles visible.

While using Toujeo, do not drive or operate heavy machinery until you know how Toujeo affects you. Don't drink alcohol or use other medicines that contain alcohol.

**The most common side effect of Toujeo is low blood sugar (hypoglycemia), which may be serious and life-threatening.** Severe hypoglycemia may cause harm to your heart or brain. Symptoms of serious low blood sugar may include shaking, sweating, fast heartbeat, and blurred vision. The long-acting effect of Toujeo may delay recovery from low blood sugar compared to shorter-acting insulins.

Please see full Important Safety Information for Toujeo throughout. Please [click here](#) or visit <https://products.sanofi.us/toujeo/toujeo.pdf> for full Prescribing Information for Toujeo.

## Important Safety Information for Toujeo U-300 (insulin glargine) injection (cont'd)

Toujeo may cause severe allergic reactions that can lead to death. Get medical help right away if you have:

- A rash over your whole body
- Shortness of breath
- Swelling of your face, tongue, or throat
- Extreme drowsiness, dizziness, or confusion
- Trouble breathing
- Fast heartbeat
- Sweating

Toujeo may have additional side effects including swelling, weight gain, low potassium, and injection site reactions which may include change in fat tissue, skin thickening, redness, swelling, and itching.

Toujeo SoloStar and Toujeo Max SoloStar are single-patient-use prefilled insulin pens. It is important to perform a safety test when using a new pen for the first time. Talk to your doctor about proper injection technique and follow instructions in the *Instruction Leaflet* that comes with your Toujeo SoloStar or Toujeo Max SoloStar pen.

**Please see full Important Safety Information for Toujeo throughout. Please [click here](https://products.sanofi.us/toujeo/toujeo.pdf) or visit <https://products.sanofi.us/toujeo/toujeo.pdf> for full Prescribing Information for Toujeo.**